



University of Crete, Department of Biology

Zebrafish Behavioral Indicators of stress & anxiety

Dr. Antonia Theodoridi
tonia.theod@gmail.com

Heraklion, May 15th, 2024

Behavioral studies in fish

There are numerous fish species used as model organisms to study fish behavior. Some of them include:

- *Danio rerio* (zebrafish)
- *Notobranchius furzeri*
- *Oryzias latipes* (medaka)
- *Poecilia reticulata* (Guppy)
- *Gasterosteus aculeatus* (three-spined stickleback)
- *Oreochromis niloticus* (Nile tilapia)



Zebrafish as a model organism

- Exhibit a wide repertoire of behaviors.
- Its small size along with the low cost housing permits high-throughput screening.
- Readily available video tracking technologies that can be coupled with zebrafish behavioral assays, providing data-rich endpoints (e.g. velocity, distance travelled, three-dimensional spatial and spatiotemporal swim path reconstructions) which are impossible to generate manually.
- Possess all major neurotransmitter systems, transporters, receptors and hormones.
- Fully sequenced genome with 70-75% of human genes having at least one zebrafish orthologue.

Zebrafish is one of the most frequently used fish species for behavioral purposes, leading to a rapid development of numerous protocols to study several behavioral phenotypes.

- Stress
- Anxiety
- Aggression
- Memory & Learning
- Reward
- Social behavior

Zebrafish as a model organism for anxiety and stress research

- Robust and easily quantifiable **cortisol** stress response.
- Clear-cut **drug-evoked phenotypes** with high predictive validity.
- Sensitivity to a wide range of **experimental stressors**, such as:

Novelty exposure	Predator exposure
Social isolation	Alarm substance
Confinement	

Disorder	Zebrafish Phenotypes
Anxiety/Fear-related behavior	<ul style="list-style-type: none">• Reduction of exploration (especially in the top part of novel environments)• Increased avoidance• Erratic behavior and freezing• Elevated cortisol and brain <i>c-fos</i>• Highly sensitive to anxiolytic and anxiogenic agents

Novelty based tests for studying anxiety

A novel environment constitutes a potentially dangerous situation for zebrafish. As a result, fish exhibit **avoidance behavior** which serves evolutionary conserved anti-predatory functions.

An animal's **exploratory behavior** in a novel environment is believed to reflect the emotional state of animals.

Tests in this category include:

1. Novel tank test (vertical exploration)
2. Open field test (horizontal exploration)
3. Light/dark preference test

Novel Tank test


A novelty based paradigm, which measures **vertical exploration**.

Zebrafish express a robust **anxiety-like response**, once introduced to the novel tank apparatus, consisting of:

- Diving to the bottom of the tank (geotaxis)
- Reduced exploration
- Increased freezing
- Erratic movements

These behavioral responses are accompanied by **physiological** responses:


- Elevated cortisol levels
- Increased breathing
- Increased heart beat frequency.



Measured parameters:

- ✓ Time spent in the top part of the tank
- ✓ Transitions to the top
- ✓ Latency to the top
- ✓ Erratic movement
- ✓ Freezing incidents
- ✓ Time spent freezing

Novel Tank test




Representative trace of the zebrafish movement in the novel tank test (6 minute trial). Different experimental manipulations show a decrease or increase in the anxious phenotype, compared to the control tank.

Novel Tank test


(A) Zebrafish strains




(B) Representative traces




Latency to upper half, s




(C) Total time in upper half, s





Total distance traveled, m



Average velocity, m/s



Light/dark preference test



Measured parameters:

- ✓ Time spent in the white zone
- ✓ Time spent in the black zone

Adult zebrafish, as well as other fish species (e.g. goldfish, guppies, minnows and tilapia), are generally believed to display an innate **aversion to bright** and white environments, and a **preference for darker** environments.


However, there are some reports which indicate a preference for the white area of the tank in zebrafish. The inconsistencies in the literature are largely attributable to differences in housing conditions (such as lighting), sex, age, social status and genetic strains.

Fish are left to acclimatize for 3-5 minutes in the centrally isolated compartment.

The separating doors are removed and fish are left to freely move in the tank for 15 minutes.

Light/dark preference test


Avoidance of the dark area & preference for the light/white area in both adult zebrafish and larvae.



Champagne *et al.*, 2010

Light/dark preference test

Avoidance of the light area & preference for the dark area in controls. Adult zebrafish exposed to the hallucinogenic drug LSD showed a preference for the light area of the tank.




Exploratory tank test


Quantification of the **exploratory** behavior of adult zebrafish.



A shoal of conspecifics is placed in the first compartment and left there undisturbed for 24 hours.



The next day the experimenter opens the “door” of the second compartment and throws some food pellets to the first and second compartment.



Fish are left free to explore the tank for 8 days. They are fed daily according to their feeding regimen.

Chronic stress protocols

Protocols that expose zebrafish to **unpredictable chronic stress** (UCS) conditions.

- The majority of them last for 14 days.
- Fish are exposed to two stressors daily. The stressors are chosen randomly every day. Some of the stressors applied in protocols of chronic stress include:

- Restraint stress
- Social isolation
- Over-crowding
- Dorsal body exposure
- Tank change
- Cold stress
- Heat stress
- Chasing
- Predator exposure
- Alarm pheromone stress

Chronic stress protocols

Behavioral, physiological and cellular responses, similar to those observed in rodents and chronically stressed humans.

More specifically, the stress protocol induced:

- Anxiety
- Cognitive impairment
- Neuroendocrine dysfunction (increased cortisol and *CRF* levels).

Impaired memory

Piato *et al.*, 2011

↑ Cortisol

Piato *et al.*, 2011; Pavlidis *et al.*, 2015

↑ CRF expression

Piato *et al.*, 2011

↓ Neurogenesis

Chakravarty *et al.*, 2013


↑ POMC, GR, MR, prolactin, BDNF, hypocretin/orexin, and c-fos expression

Pavlidis *et al.*, 2015



Aggression in zebrafish

Zebrafish in nature form shoals, yet when put in pairs they often exhibit an aggressive behavior that leads to the establishment of a social hierarchy. The “winner” of this dyadic interaction is considered the **dominant** individual, while the “loser” is the **subordinate**.

The established hierarchy consists a form of **social stress** for both the dominant and the subordinate, therefore, zebrafish exhibit anxiety-like behavior. Both dominants and subordinates show higher cortisol levels compared to controls (Pavlidis *et al.*, 2011).



Aggressive behavior in zebrafish



Oliveira *et al.*, 2011

Displaying consists of an approach to the conspecific followed by a turn to the left or right with fins erect.

Circling is considered another form of lateral display that may last an extended period and during which the fish rise in the water column.

Chasing is a pursuit or a quick approach to an individual.

Bites consist of closing the mouth against another individual.

Freezing is the term that describes the immobility state with retracted fins.

A **fleeing** individual moves away from a pursuing individual for an extended period of time.


Dominant behavior consists mainly of **chasing** and **biting**, while subordinate behavior is expressed as **fleeing** and **freezing**.

Unconditioned social interaction test / Paired aggression test



Pairs of same sex adult zebrafish are introduced in 2L tanks

2 h →



Recording of the pair's behavior for 5 minutes.


→

Behavioral analysis and quantification of the **dominant** and **subordinate** behavior




- Number of attacks from the dominant
- Total duration of chasing
- Total duration of subordinate's freezing behavior

A. Dominant Behavior




Acute exposure to fluoxetine can alter the aggressive behavior of adult male zebrafish.

B. Subordinate Behavior



Aggressive behaviors in zebrafish



- a. **Mirror-elicited fight:** The arena is divided by a partition which contains one mirror in each side. A second partition is added on each side in order to cover the mirror. To elicit the mirror fight, the outer partitions are removed and the fish are allowed to interact with their own reflection.
- b. **Real opponent fight:** The arena is divided by a removable opaque partition. Once the partition is removed, the two same-sex conspecifics are allowed to interact.


Mirror test

An established protocol to study **social** and **aggressive** behavior in adult zebrafish.


Not possible to observe the full repertoire of aggressive behavior and it has been shown that zebrafish have different endocrinological and genetic responses to a mirror “fight” (Teles *et al.*, 2013), but the risk of injury is avoided.

Measured parameters:

- ✓ Frequency of attacking the mirror
- ✓ Total duration of biting the mirror
- ✓ Number of contacts with the mirror
- ✓ Latency of approaching the mirror



Mirror test




Norton *et al.*, 2011



Representative trace of the swimming behavior of WT (b) and mutants with a robust aggressive phenotype (knock-out of **fibroblast growth factor receptor 1a** (Fgfr1a)) (c).

Tests assessing social behavior

Shoal cohesion test




Social preference test




Tests assessing social behavior

A


Conspecific Target fish Empty zone




Group Target fish Conspecific



Non-kin Target fish Kin




Unfamiliar Target fish Familiar




B

Normal school



Disrupted, loose school



Predator avoidance test

Zebrafish natural predators include among others Indian leaf fish (*Nandus nandus*), freshwater garfish (*Xenentodon* spp.), catfish (*Mystus bleekeri*), as well as avian predators.

In laboratory conditions, the presence of the Indian leaf fish induces fear-like responses in zebrafish, which consist of:

- Increased frequency of escape efforts and
- erratic movements post exposure to a predator (Bass & Gerlai, 2008).

Depending on the species of predator that they are exposed to, zebrafish adjust their defensive and fear-like behaviors (Ahmed *et al.*, 2012).



Kalueff *et al.*, 2014



Indian leaf fish, *Nandus nandus*

Predator avoidance test


Exposure to both live predator and the robotic model induced a robust avoidance response in zebrafish. Contrary to this, computer-animated images failed to elicit a fear-like behavioral response.





(a) Live red tiger oscar; (b) robotic predator designed after the live oscar; and (c) snapshot of the computer-animated of live Oscar

T-maze

- Learning capacity
- Long-term memory
- Short-term memory
- Memory plasticity




T-maze test



T-maze

Alternative structure

One arm has an enriched environment which is favorable to zebrafish (plants and marbles).



Assessing stress response in zebrafish larvae

Fewer behavioral protocols have been established for zebrafish larvae. The most commonly used are:

- Acute stress response to a swirling stressor
- Tapping assay
- Light/Dark preference
- Open field test

Behavioral studies in zebrafish larvae

Adult zebrafish used for reproduction





Fertilised eggs



Larvae 5 days post fertilization (dpf)




Transfer of larvae in cell culture plates



Observation Chamber

Assessing stress response in zebrafish larvae




Larvae with different genetic manipulations related to genes regulating the HPI axis display differences in light/dark preference and anxiety levels (thigmotaxis).

Behavioral analysis software packages

Noldus



Ethovision



Daniovision

Behavioral analysis software packages

Viewpoint

Zebrabox




Zebracube




Behavioral analysis software packages

Smart 3.0

Y-maze Spontaneous Alternation



Zebrafish Larvae Activity



Open-source behavioral analysis software packages

<i>Software</i>	<i>Operative system/program</i>	<i>Types of analyses</i>	<i>Types of input</i>	<i>Types of output</i>	<i>Webpage</i>
wrMTrck	Windows and Mac OS X/ JAVA-ImageJ	Total length, average speed, area, perimeter, and trajectories.	AVI files with jpg compression	txt, xls, tiff files and AVI videos	www.phage.dk/plugins/ wrmtrck.html
Mouse Behavior Tracker	Windows, Mac OS X and Linux/JAVA-ImageJ	Distance and average velocity.	AVI or MPEG-compressed AVI files, Mp4	Txt or xls files	www.BioTechniques.com/ article/114607.
AnimalTracker	Windows, Mac OS X and Linux/JAVA-ImageJ	Total length, average speed, and time spent in ROI.	AVI files with jpg compression	txt, xls, tiff files and AVI videos	animaltracker.elte.hu
idTracker	Windows/MATLAB	Trajectories, identification of one animal in different videos and ROI.	Compatible with MATLAB, uncompressed AVI or MPEG-compressed AVI files	X and Y coordinates and images files	www.idtracker.es
Mousetracker	Windows, Linux and Mac OS X/Pascal-Delphi-MS Excel	Velocity, acceleration, and time spent in ROI.	AVI format	XY coordinates can be copied directly	www.neuro.ufrn.br/ software/mouselabtracker
JAABA	Windows, Mac OS X and Linux/MATLAB	Bites, persecution, sexual behavior, angle of turn, grooming, jump, walk, immobilization, and touch. Locomotion and ROI.	Several formats and resolutions. X and Y coordinates	MATLAB files	http://jaaba.sourceforge.net https://www.janelia .org/lab/branson-lab
Ctrax	Windows and Mac OS X/ Phyton—MATLAB	Trajectories, velocities, speed, position, and turning speed histograms.	Common digital video formats, mainly AVI	csv and mat files. Converts the file to .ann extension	ctrax.sourceforge.net
VideoHacking	Windows, Mac OS X and Linux/Phyton—Open CV	Velocity, acceleration, total length, average speed, and time spent in ROI.	Common digital video formats	Graphical interface to view data summary	faculty.ithaca.edu/iwoods/ docs/
ToxTrack/ToxId	Windows/C++	Total distance, speed, acceleration, time near the walls (measure of anxiety), and ROI.	AVI or MPEG-compressed AVI files	txt, xls, tiff files and AVI videos	https://sourceforge .net/projects/toxtrac/
EthoWatcher	Windows/C++	Frequency, duration, and latency of each behavior.	AVI or MPEG-compressed AVI files	csv files	http://ethowatcher .paginas.ufsc.br
MouseMove	Windows/LabView—ImageJ	Distance, average velocity, acceleration, curvature, stationary fraction, laterality y ROI.	AVI or MPEG-compressed AVI files	csv files	https://www.nature.com/ articles/srep16171#s3, Supplementary File 2
Cowlog	Windows, Mac OS X and Linux/Java—html	Analysis of different behaviors can be set (tapping a button when the event occurs)	Common digital video formats	csv files	cowlog.org

AVI, audio video interleaved; MJPEG, motion joint photographic experts group; ROI, region of interest.

Open-source behavioral analysis software packages

Review Article | [Published: 29 July 2021](#)

A review of 28 free animal-tracking software applications: current features and limitations

[Veronica Panadeiro](#), [Alvaro Rodriguez](#) , [Jason Henry](#), [Donald Wlodkowic](#) & [Magnus Andersson](#)

[Lab Animal](#) **50**, 246–254 (2021) | [Cite this article](#)

Deep-learning methods in behavioral analyses


www.nature.com/npp

Neuropsychopharmacology



ARTICLE **OPEN**

Deep learning-based behavioral analysis reaches human accuracy and is capable of outperforming commercial solutions

Oliver Sturman^{1,2}, Lukas von Ziegler^{1,2}, Christa Schläppi^{1,2}, Furkan Akyol^{1,2}, Mattia Privitera^{1,2}, Daria Slominski^{1,2}, Christina Grimm^{2,3}, Laetitia Thieren^{2,4}, Valerio Zerbi^{2,3}, Benjamin Grewe^{2,5,6} and Johannes Bohacek ^{1,2}

Deep-learning methods in behavioral analyses

nature
neuroscience

TECHNICAL REPORT

<https://doi.org/10.1038/s41593-018-0209-y>

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning

Alexander Mathis ^{1,2}, Pranav Mamidanna¹, Kevin M. Cury³, Taiga Abe³, Venkatesh N. Murthy ², Mackenzie Weygandt Mathis ^{1,4,8*} and Matthias Bethge^{1,5,6,7,8}



University of
St Andrews

[Home](#) [Profiles](#) [Research output](#) [Datasets/Software](#) [Research units](#) [Projects](#) [Activities](#) [...](#)

PoseR - A deep learning toolbox for decoding animal behavior

Pierce Mullen, Maarten Frans Zwart

School of Psychology and Neuroscience, Centre for Biophotonics, Institute of Behavioural and Neural Sciences

DeepLabCut in zebrafish behavioral studies


© 2022. Published by The Company of Biologists Ltd | Biology Open (2022) 11, bio059443. doi:10.1242/bio.059443



RESEARCH ARTICLE


Beyond bold versus shy: Zebrafish exploratory behavior falls into several behavioral clusters and is influenced by strain and sex

Neha Rajput, Kush Parikh and Justin W. Kenney*



- Over 400 zebrafish
- 4 strains (AB, TL, TU, WIK)
- Both sexes
- 3D swim traces

4 behavioral types





DeepLabCut in zebrafish behavioral studies

Pose analysis in free-swimming adult zebrafish, *Danio rerio*:
“fishy” origins of movement design

Jagmeet S. Kanwal^{1*}, Bhavjeet Sanghera^{†1,2}, Riya Dabbi¹ and Eric Glasgow³

¹Department of Neurology, Georgetown University Medical Center, Washington, DC, USA, ²University of Miami, Coral Gables, FL, USA ³Department of Oncology, Georgetown University Medical Center, USA



References

- Kalueff, A.V., Stewart, A.M., Gerlai, R. (2014). Zebrafish as an emerging model for studying complex brain disorders. *Trends Pharmacol Sci* 35(2): 63-75.
- Cachat, J. et al. (2010). Measuring behavioral and endocrine responses to novelty stress in adult zebrafish. *Nature protocols*, 5(11), 1786–1799.
- Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F., Elkhayat, S.I., Bartels, B.K., Tien, A.K., Tien, D.H., et al. (2009). Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behavioural brain research*, 205(1), 38–44.
- Piato, Â.L. et al. (2011). Unpredictable chronic stress model in zebrafish (*Danio rerio*): behavioral and physiological responses. *Progress in neuro-psychopharmacology & biological psychiatry*, 35(2), 561–7.
- Pavlidis, M., Theodoridi, A. and Tsalafouta, A. (2015). Neuroendocrine regulation of the stress response in adult zebrafish, *Danio rerio*. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 60, 121–131.
- Chakravarty, S. et al. (2013). Chronic unpredictable stress (CUS)-induced anxiety and related mood disorders in a zebrafish model: altered brain proteome profile implicates mitochondrial dysfunction. *PloS one*, 8(5), e63302.
- Pavlidis, M. et al. (2011). Adaptive changes in zebrafish brain in dominant-subordinate behavioral context. *Behavioural brain research*, 225(2), 529–37.
- Oliveira, R.F., Silva, J.F. and Simões, J.M. (2011). Fighting Zebrafish: Characterization of Aggressive Behavior and Winner–Loser Effects. *Zebrafish*, 8(2), 73–81.
- Theodoridi, A., Tsalafouta, A., Pavlidis, M. (2017). Acute Exposure to Fluoxetine Alters Aggressive Behavior of Zebrafish and Expression of Genes Involved in Serotonergic System Regulation. *Front Neurosci*, 11, 223.
- Nunes, A., Ruhl, N., Winberg, S. & Oliveira, R. (2017). Social Phenotypes in Zebrafish. *The rights and wrongs of Zebrafish: Behavioral Phenotyping of Zebrafish*.
- Teles, M.C. et al. (2013). Social modulation of brain monoamine levels in zebrafish. *Behavioural brain research*, 253C, 17–24.
- Norton, W.H.J. et al. (2011). Modulation of *Fgfr1a* signaling in zebrafish reveals a genetic basis for the aggression–boldness syndrome. *Journal of Neuroscience*, 31(39): 13796–13807

References

- Stewart, A.M., Nguyen, M., et al. (2014). Developing zebrafish models of autism spectrum disorder (ASD). *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 50, 27–36.
- Bass SL, Gerlai R. (2008). Zebrafish (*Danio rerio*) responds differentially to stimulus fish: the effects of sympatric and allopatric predators and harmless fish. *Behav Brain Res*, 186(1):107-17.
- Ahmed O, Seguin D, Gerlai R. (2011). An automated predator avoidance task in zebrafish. *Behav Brain Res*, 216(1): 166-71.
- Fabrizio, L., Violet, M., Jasmine, L., Simone, M., Maurizio, P. (2015). Acute caffeine administration affects zebrafish response to a robotic stimulus. *Behavioural Brain Research*, 289, 48-54.
- Faught, E. and Vijayan, M.M. (2018). The mineralocorticoid receptor is essential for stress axis regulation in zebrafish larvae. *Scientific Reports*, 8(1), 1–11.
- Franco-Restrepo, J.E., Forero, D.A., Vargas, R.A. (2019). A Review of Freely Available, Open-Source Software for the Automated Analysis of the Behavior of Adult Zebrafish. *Zebrafish*, 16:3, 223-232.
- Panadeiro, V., Rodriguez, A., Henry, J., Wlodkowic, D., Andersson, M. (2021). A review of 28 free animal-tracking software applications: current features and limitations. *Lab Anim (NY)*, 50(9):246-254.
- Sturman, O., von Ziegler, L., Schläppi, C. et al. (2020). Deep learning-based behavioral analysis reaches human accuracy and is capable of outperforming commercial solutions. *Neuropsychopharmacol.* 45, 1942–1952.
- Mullen, P., & Zwart, M. F. (2023). PoseR - A deep learning toolbox for decoding animal behavior. *bioRxiv*.
- Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience*.
- Rajput, N., Parikh, K., Kenney, J.W. (2022). Beyond bold versus shy: Zebrafish exploratory behavior falls into several behavioral clusters and is influenced by strain and sex. *Biol Open*; 11(8):bio059443.
- Jagmeet S. Kanwal, Bhavjeet Sanghera, Riya Dabbi, Eric Glasgow. (2023). Pose analysis in free-swimming adult zebrafish, *Danio rerio*: “fishy” origins of movement design. *bioRxiv*.

Thank you!

Any questions?
tonia.theod@gmail.com

